

Exclusive $B \rightarrow K^{**}\gamma$ Decays in the QCD LCSR approach*

A. S. Safir

DESY, Deutsches Elektronen-Synchrotron, D-22603 Hamburg, Germany
 E-mail : safir@mail.desy.de

February 1, 2008

Abstract

We predict contributions of higher K -resonances to the radiative rare decays $b \rightarrow s\gamma$, in the framework of the QCD sum rules on the light cone (LCSR). Our calculations are restricted to the leading twist-two operators for $K^*(892)$ and to the asymptotic wave function for the other K^{**} -mesons. Using experimental data on the semileptonic $\tau \rightarrow K^{**}\nu_\tau$ decays, we extract the corresponding decay constants for vector and axial-vector K^{**} -mesons. We present results for the corresponding branching ratios and compare them with the existing theoretical predictions.

1 Introduction

The study of radiative decays based on the flavour-changing neutral $b \rightarrow (s, d) + \gamma$ current transition is of crucial importance for testing the flavour sector of the Standard Model and probing for new physics. In the standard model, the short distance contribution to rare B-decays is dominated by the top quark, and long distance contributions by form factors. Precise measurements of these transition will not only provide a good estimate of the top quark mass and the CKM matrix elements V_{td} , V_{ts} , V_{tb} , but also of the hadronic properties of B -mesons, namely form factors which in turn would provide a good knowledge of the corresponding dynamics and more hint for the non-perturbative regime of QCD.

Experimental measurements of exclusive $B \rightarrow K^*\gamma$ branching ratios have been reported by the BABAR, CLEO and BELLE Collaborations, with the results:

$$10^5 \text{Br}(\bar{B}^0 \rightarrow \bar{K}^{*0}\gamma) = \begin{cases} 4.23 \pm 0.40 \pm 0.22 & [1] \\ 4.55^{+0.72}_{-0.68} \pm 0.34 & [2] \\ 4.96 \pm 0.67 \pm 0.45 & [3] \end{cases}$$

and also of the inclusive rate [2–4]:

$$10^4 \text{Br}(B \rightarrow X_s\gamma) = (3.22 \pm 0.40)$$

*Talk presented at the XXXVIIth Rencontres de Moriond: Electroweak Interactions and Unified Theories, Les Arcs, France, 9-16 March 2002.

However, the first observation of the rare B -decay to the orbitally excited strange mesons has been reported by CLEO [2], and recently by BELLE [3]:

$$10^5 Br(B \rightarrow K_2^*(1430)\gamma) = \begin{cases} (1.66_{-0.53}^{+0.59} \pm 0.13) & [2] \\ (1.26 \pm 0.66 \pm 0.10) & [3] \end{cases}$$

These important experimental measurements provide a crucial challenge to the theory. Whereas many theoretical approaches have been employed to predict the exclusive $B \rightarrow K^*(892)\gamma$ decay rate, less attention has been devoted to rare radiative B -decays to excited strange mesons [5–8]. Most of these theoretical approaches rely on non-relativistic quark models [5, 6], HQET [7] and relativistic model [8]. However there is a large spread between different results, due to different treatments of the long distance effects.

In this talk we present the results of [9], where a systematic analysis of the electromagnetic penguin form factor $F_1^{K^{**}}(0)$ governing the exclusive rare B -decays to orbitally excited K^{**} -mesons was performed in the framework of the QCD sum rules on the light cone [10].

2 General framework

At the quark level, the rare semileptonic decay $b \rightarrow s\gamma$ can be described in terms of the effective Hamiltonian obtained by integrating out the top quark and $W^\pm$ bosons:

$$H_{eff} = -4 \frac{G_F}{\sqrt{2}} V_{ts}^* V_{tb} \sum_{i=1}^8 C_i(\mu) O_i(\mu) . \quad (1)$$

where V_{ij} are the corresponding CKM matrix elements and G_F is the Fermi coupling constant. Following the notation and the convention of ref. [11], regarding the operator basis, one can test the model dependence of the form factors for the exclusive decay in the ratio of the exclusive-to-inclusive radiative decay branching ratio:

$$R_{K^{**}} \equiv \frac{BR(B \rightarrow K^{**} \gamma)}{BR(B \rightarrow X_s \gamma)} \simeq F_1^{K^{**}}(0)^2 \zeta(m_s, m_b, m_{K^{**}}, \dots) \quad (2)$$

where $\zeta(m_s, m_b, m_{K^{**}}, \dots)$ is a kinematic function which can be found in [9]. With this normalization, one eliminates the uncertainties from the CKM matrix elements and the short distance contribution. Thus, we are left in (2) with unknown form factors $F_1^F(0)$, which we will derive using QCD sum rules on the light cone.

The starting point of our sum rule is to consider the correlation function

$$i \int dx e^{iqx} \times \langle K^{**}(p, \epsilon) | T \{ \bar{\psi}(x) \sigma_{\mu\nu} (\sigma_{\mu\nu} \gamma_5) q^\nu b(x) \bar{b}(0) i \gamma_5 \psi(0) \} | 0 \rangle \quad (3)$$

Hereafter we use ψ as a generic notation for the field of the light quark. The hadronic representation of (3) is obtained by inserting a complete set of states including the B -meson ground state, higher resonances and the non-resonant states with B -meson quantum numbers. After writing down the dispersion relation in $(p+q)^2$, we can separate the contribution of the B -meson as the pole contribution.

The possibility to calculate the correlator (3) in the region of large space-like momenta $(p+q)^2 < 0$ is based on the expansion of the T -product of the currents in (3) near the light-cone $x^2 = 0$ which is expressed through matrix elements of non-local operators, sandwiched in between the

Table 1: *Central values of the pseudoscalar, vector, scalar and axial vector K^{**} -meson decay constants (in MeV) and the corresponding form factors.*

	$K^*(892)$	$K_1^*(1270)$	$K_1(1400)$	$K^*(1410)$	$K_0^*(1430)$	$K_1(1650)$	$K^*(1680)$
J^P	1^-	1^+	1^+	1^-	0^+	1^+	1^-
f_i	210	122	91	86	79	86	86
$F_1^{K^{**}}(0)$	$0.32^{+0.06}_{-0.06}$	$0.14^{+0.03}_{-0.03}$	$0.098^{+0.02}_{-0.02}$	$0.094^{+0.02}_{-0.02}$		$0.091^{+0.02}_{-0.02}$	$0.091^{+0.02}_{-0.02}$

Table 2: *Comparison of our results for the ratio $R_F[\%]$ with previous works.*

Meson	$R_F[\%]$				
	ref. [9]	ref. [7]	ref. [6]	ref. [5]	ref. [8]
K	forb.				
$K^*(892)$	$10.0^{+4.0}_{-4.0}$	16.8 ± 6.4	$3.5 - 12.2$	4.5	15^{+3}_{-3}
$K^*(1430)$	forb.				
$K_1(1270)$	$2.0^{+0.8}_{-0.8}$	$4.3^{+1.6}_{-1.6}$	$4.5 - 10.1$	forb./6.0	$1.5^{+0.5}_{-0.5}$
$K_1(1400)$	$0.9^{+0.4}_{-0.4}$	$2.1^{+0.9}_{-0.9}$	$6.0 - 13.0$	forb./6.0	$2.6^{+0.6}_{-0.6}$
$K_2^*(1430)$	$5.0^{+2.0}_{-2.0}$	$6.2^{+2.9}_{-2.9}$	$17.3 - 37.1$	6.0	$5.7^{+1.2}_{-1.2}$
$K^*(1680)$	$0.7^{+0.3}_{-0.3}$	$0.5^{+0.2}_{-0.2}$	$1.0 - 1.5$	0.9	
$K_2(1580)$		$1.7^{+0.4}_{-0.4}$	$4.5 - 6.4$	4.4	
$K(1460)$	forb.				
$K^*(1410)$	$0.8^{+0.4}_{-0.4}$	$4.1^{+0.6}_{-0.6}$	$7.2 - 10.6$	7.3	
$K_0^*(1950)$	forb.				
$K_1(1650)$	$0.8^{+0.3}_{-0.3}$	$1.7^{+0.6}_{-0.6}$	not given	not given	

vacuum and the meson state. These matrix elements define the light-cone meson wave functions. We restricted our calculations to the leading twist-2 operator for the $K^*(892)$, as in [12], and to the asymptotic wave function for the other K^{**} -mesons. The latter choice is simply based on the fact that using QCD sum rules, it is impossible to get rid of the lower-lying states contributions from these higher resonances.

However, nothing is known about the corresponding K^{**} -decay constants, and one has to predict them. For that, we have used recent data [13] on semileptonic $\tau \rightarrow (K^{**})\nu_\tau$ decays * to obtain them [9]. We present in table 1 the corresponding decay constants. For $K_2^*(1430)$, we have constrained the corresponding decay constant with the recent data [2] on $B \rightarrow K_2^*(1430)\gamma$.

Following the basic steps of the QCD sum rules on the light-cone, as described above, and using the experimental K^{**} -decay constants, we show in table 1 the corresponding form factors. Finally, in table 2 we compare our results for the ratio $R_{K^{**}}[\%]$ with previous works [5–8].

3 Summary

Motivated by the first observation of the radiative decay $B \rightarrow K_2^*(1430)\gamma$, we have investigated rare radiative B decays to orbitally excited K^{**} -mesons. First, we have presented an alternative method of calculating the transition form factors and related decays using the QCD sum rules on the light-cone. For that, We have extracted the unknown K^{**} -decay constants using the recent

*These quantities contribute to the transition rates for pseudoscalar, vector, scalar and axial vector emission.

data [13] on semileptonic $\tau \rightarrow K^{**}\nu_\tau$ decays.

For $K_2^*(1430)$, we have constrained the corresponding decay constant with the recent data [2] on $B \rightarrow K_2^*(1430)\gamma$. We find that if $f_{K_2^*(1430)} = (140 - 180) \text{ MeV}$, a substantial fraction (3.0 – 7.0)% of the inclusive $b \rightarrow s\gamma$ branching ratio goes into the $K_2^*(1430)$ channel, in a good agreement with recent CLEO data [2]. Our prediction for the $B \rightarrow K^*(892)\gamma$ branching fraction yields to (6.0 – 14.0)%, also in good agreement with the experimental data [1–3].

In order to make comparison of our results with previous calculations, we have tabulated our results together with results of [5], [6], [7] and [8] in table 2. As far as decays into higher K -resonances are concerned, our results are in general in much better agreement with [7] than [5] and [6], apart from the $K^*(1410)$ -channel where the difference is more significant. Finally, it should be noticed that the theoretical uncertainties in our light-cone sum rules are the wave functions and the decay constants of the K^{**} -mesons. The accuracy of our calculation can be substantially improved by taking into account the wave functions of twist-3 and twist-4 for the $B \rightarrow K^*(892)\gamma$ decay, and going beyond the asymptotic form for the other decay modes. To reduce the uncertainties on the K^{**} -mesons decay constants, one can determine them independently using QCD sum rules for the two-point correlator of the corresponding currents.

Acknowledgments

I would like to thank the Organizers of the XXXVIIth Rencontres de Moriond for their financial support. I express my gratitude to A. Ali for critical reading of the manuscript.

References

- [1] BABAR Collaboration, B. Aubert et al., Phys. Rev. Lett. **88**, 101805 (2002).
- [2] CLEO Collaboration, Phys. Rev. Lett. **84** (2000) 5283; hep-ex/9908022; hep-ex/0108032.
- [3] BELLE Collaboration, hep-ex/0104045; hep-ex/0107065.
- [4] ALEPH Collaboration, R. Barate et al., Phys. Lett. **B429** (1998) 169.
- [5] T. Altomari, Phys. Rev. D **37**, 677 (1988).
- [6] A. Ali et al., Phys. Lett. **B298**, 195 (1993).
- [7] S. Veseli and M.G. Olsson, Phys.Lett.**B367**(1996)309.
- [8] D. Ebert et al., Phys. Lett. **B495** (2000) 309; Phys.Rev.**D64** (2001) 054001.
- [9] A. S. Safir, Eur. Phys. J. directC **15**, 1 (2001).
- [10] I. I. Balitsky et al., Sov. J. Nucl. Phys. **44** (1986) 1028; Nucl. Phys.**B312** (1989) 509; V. M. Braun and I. E. Filyanov, Z. Phys. **C48** (1990) 239; V. L. Chernyak, A. R. Zhitnisky, Phys. Rep. **112** (1984) 173; A. S.Gorsky Sov. J. Nucl. Phys. **41** (1985) 1008; ibid. **45** (1987) 512.
- [11] A. Ali and C. Greub, Z. Phys. **C49** (1991) 431; A. J. Buras et al. Nucl. Phys. **B424** (1994) 374; K. Chetyrkin et al., Phys. Lett. **B400** (1997) 206; Erratum-ibid **B425** (1998) 414.
- [12] A. Ali, V. M. Braun and H. Simma, Z. Phys. **C67** (1994) 437.
- [13] Particle Data Group, Lepton Summary Table, Eur. Phys. J. **C15** (2000) 23.